



Centrum voor Wiskunde en Informatica

**REPORTRAPPORT**

A Godunov-type scheme with applications in hydrodynamics

E.H. van Brummelen

Modelling, Analysis and Simulation (MAS)

**MAS-R9829 December 1998**

Report MAS-R9829  
ISSN 1386-3703

CWI  
P.O. Box 94079  
1090 GB Amsterdam  
The Netherlands

CWI is the National Research Institute for Mathematics and Computer Science. CWI is part of the Stichting Mathematisch Centrum (SMC), the Dutch foundation for promotion of mathematics and computer science and their applications.

SMC is sponsored by the Netherlands Organization for Scientific Research (NWO). CWI is a member of ERCIM, the European Research Consortium for Informatics and Mathematics.

Copyright © Stichting Mathematisch Centrum  
P.O. Box 94079, 1090 GB Amsterdam (NL)  
Kruislaan 413, 1098 SJ Amsterdam (NL)  
Telephone +31 20 592 9333  
Telefax +31 20 592 4199

# A Godunov-Type Scheme with Applications in Hydrodynamics

E.H. van Brummelen

CWI

*P.O. Box 94079, 1090 GB Amsterdam, The Netherlands*

## ABSTRACT

In spite of the absence of shock waves in most hydrodynamic applications, sufficient reason remains to employ Godunov-type schemes in this field. In the instance of two-phase flow, the shock capturing ability of these schemes may serve to maintain robustness and accuracy at the interface. Moreover, approximate Riemann solvers have greatly relieved the initial drawback of computational expensiveness of Godunov-type schemes. In the present work we develop an Osher-type flux-difference splitting approximate Riemann solver and we examine its application in hydrodynamics. Actual computations are left to future research.

*1991 Mathematics Subject Classification:* 35L65, 35L67, 76L05, 73D20, 76T05.

*Keywords and Phrases:* hydrodynamics, Godunov-type schemes, hyperbolic conservation laws, two-phase flows, surface waves, interface capturing.

*Note:* This work was performed under a research contract with the Maritime Research Institute Netherlands and was carried out under CWI-project MAS2.1 "Computational Fluid Dynamics".

## 1. INTRODUCTION

The advantages of Godunov-type schemes [God59] in hydrodynamic flow computations are not as widely appreciated as in gas dynamics applications. Admittedly, the absence of supersonic speeds and hence shock waves in incompressible flow (the prevailing fluid model in hydrodynamics) reduces the necessity of advanced shock capturing schemes. Nevertheless, many reasons remain to apply Godunov-type schemes in hydrodynamics: Firstly, these schemes have favourable robustness properties due to the inherent upwind treatment of the flow. Secondly, they feature a consistent treatment of boundary conditions. Thirdly, (higher-order accurate) Godunov-type schemes display low dissipative errors, which is imperative for an accurate resolution of boundary layers in viscous flow. Finally, the implementation of these schemes in conjunction with higher-order limited interpolation methods, to maintain accuracy and prevent oscillations in regions where large gradients occur (see, e.g., [Swe84, Spe87]), is relatively straightforward.

In addition, Godunov-type schemes can be particularly useful in hydrodynamics in case of two-phase flows, e.g., flows suffering cavitation and free surface flows. In these situations, an interface exists between the primary phase (water) and the secondary phase (air, damp, etc.) and fluid properties may vary discontinuously across the interface. In our opinion, the ability of Godunov-type schemes to capture discontinuities is then very useful to maintain robustness and accuracy at the interface. Examples of such interface capturing can be found in, for instance, [MOS92, CHMO96, KP97].

A disadvantage of the method originally proposed by Godunov is that it requires the solution of an associated Riemann problem with each flux evaluation. In practice, many such evaluations are performed during an actual computation. Consequently, the method is notorious for its high computational costs. To relieve this problem, several approaches have been suggested to reduce the computational costs of the flux evaluations involved, by approximating the Riemann solution. Examples of such approximate Riemann solvers are the flux vector splitting schemes (such as those

of Van Leer [Lee82] and Steger & Warming [SW81]) and the flux difference splitting schemes (such as Roe's [Roe81] and Osher's [OS82]).

In the present work we develop an Osher-type flux-difference splitting scheme for the approximate solution of the Riemann problem and we investigate its application in hydrodynamics. Details are presented for the Euler equations for three types of fluids that are used to model the behaviour of water, viz., a genuinely compressible fluid, an artificially compressible fluid and a genuinely incompressible fluid, and for a two-phase flow. As a preliminary, we examine the Riemann problem. Next, we give an outline of Osher's approximate Riemann solver. Analysis shows that Osher's scheme suffers loss of accuracy in the presence of centered shock waves and therefore a modified scheme is proposed. Finally, we present the specifics for the aforementioned hydrodynamic applications. Actual computations are deferred to future research.

## 2. RIEMANN PROBLEM

In this section we investigate the Riemann Problem:

**Definition 2.1** Let  $\mathbf{q} \in \mathbb{R}^n = (q_1, \dots, q_n)^T$ ,  $(x, t) \in \mathbb{R} \times \mathbb{R}^+$  and  $\mathbf{f} \in C^1(\mathbb{R}^n, \mathbb{R}^n)$ . Consider the Cauchy problem

$$\partial_t \mathbf{q} + \partial_x \mathbf{f}(\mathbf{q}) = 0, \quad \forall x \in \mathbb{R}, t \in \mathbb{R}^+, \quad (2.1a)$$

subject to the initial condition

$$\mathbf{q}(x, 0) = \begin{cases} \mathbf{q}_L, & \text{if } x < 0, \\ \mathbf{q}_R, & \text{if } x > 0, \end{cases} \quad (2.1b)$$

with  $\mathbf{q}_L$  and  $\mathbf{q}_R$  constant. The initial value problem (2.1a) and the initial condition (2.1b) define the Riemann problem.

First, an introductory analysis is presented. Subsequently, we obtain the general solution to (2.1).

### 2.1 Preliminary Analysis

Let  $\mathbf{A}(\mathbf{q})$  denote the Jacobian of  $\mathbf{f}(\mathbf{q})$ ,  $\mathbf{A}(\mathbf{q}) = \partial_{\mathbf{q}} \mathbf{f}(\mathbf{q})$ , and let  $\lambda_k(\mathbf{q})$ ,  $k = 1, 2, \dots, n$ ,  $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$ , be its eigenvalues and  $\mathbf{r}_k(\mathbf{q})$  the corresponding eigenvectors. Equation (2.1a) constitutes a hyperbolic system if the eigenvalues  $\lambda_k(\mathbf{q})$  are real and nonzero. Then, the matrix  $\mathbf{A}(\mathbf{q})$  can be decomposed with respect to a basis of its eigenvectors:

$$\mathbf{A}(\mathbf{q}) = \mathbf{R}(\mathbf{q}) \cdot \mathbf{\Lambda}(\mathbf{q}) \cdot \mathbf{R}(\mathbf{q})^{-1}, \quad (2.2)$$

where  $\mathbf{\Lambda}(\mathbf{q}) = \text{diag}(\lambda_1(\mathbf{q}), \dots, \lambda_n(\mathbf{q}))$  and the matrix  $\mathbf{R}(\mathbf{q}) = (\mathbf{r}_1(\mathbf{q}), \dots, \mathbf{r}_n(\mathbf{q}))$  contains the eigenvectors. From [Lax57] we adopt the following classification of the eigenpairs  $(\lambda_k(\mathbf{q}), \mathbf{r}_k(\mathbf{q}))$ :

**Definition 2.2** Consider the matrix  $\mathbf{A}(\mathbf{q}) \in \mathbb{R}^{n \times n}$ . Let  $\lambda_k(\mathbf{q})$ ,  $k = 1, 2, \dots, n$ , be its eigenvalues and  $\mathbf{r}_k(\mathbf{q})$  the corresponding eigenvectors. An eigenvalue  $\lambda_k(\mathbf{q})$  and an eigenvector  $\mathbf{r}_k(\mathbf{q})$  are called genuinely nonlinear on a subdomain  $\Omega \subseteq \mathbb{R}^n$  if

$$\partial_{\mathbf{q}} \lambda_k(\mathbf{q}) \cdot \mathbf{r}_k(\mathbf{q}) \neq 0, \quad \forall \mathbf{q} \in \Omega. \quad (2.3)$$

An eigenvalue  $\lambda_k(\mathbf{q})$  and an eigenvector  $\mathbf{r}_k(\mathbf{q})$  are said to be linearly degenerate on  $\Omega$  if

$$\partial_{\mathbf{q}} \lambda_k(\mathbf{q}) \cdot \mathbf{r}_k(\mathbf{q}) = 0, \quad \forall \mathbf{q} \in \Omega. \quad (2.4)$$

The eigenvalues that are genuinely nonlinear for all  $\mathbf{q} \in \mathbb{R}^n$  are related to rarefaction waves and shocks in the solution of the Riemann problem. The eigenvalues that are linearly degenerate on  $\mathbb{R}^n$  correspond to contact discontinuities in the solution. More complex contact phenomena can occur for eigenvalues that are neither genuinely nonlinear nor linearly degenerate on  $\mathbb{R}^n$ , see, e.g., [LeV90, pages 48–50].

With each of the eigenpairs  $(\lambda_k(\mathbf{q}), \mathbf{r}_k(\mathbf{q}))$  we associate two paths in state space. Firstly, the  $k$ -shock path:

**Definition 2.3** Consider the hyperbolic system (2.1a). The  $k$ -shock path through  $\mathbf{q}_L$  is the set

$$\mathcal{S}_k(\mathbf{q}_L) = \{\mathbf{q} \in \mathbb{R}^n \mid s(\mathbf{q}; \mathbf{q}_L)(\mathbf{q} - \mathbf{q}_L) = \mathbf{f}(\mathbf{q}) - \mathbf{f}(\mathbf{q}_L)\}, \quad (2.5)$$

where  $s(\mathbf{q}; \mathbf{q}_L)$  is referred to as the shock speed.

Secondly, we distinguish the  $k$ -path:

**Definition 2.4** Consider the hyperbolic system (2.1a). The  $k$ -path through  $\mathbf{q}_L$  is the set

$$\mathcal{R}_k(\mathbf{q}_L) = \{\mathbf{q} \in \mathbb{R}^n \mid \mathbf{q} = \mathbf{h}(\xi), \xi \in \mathbb{R}\} \quad (2.6)$$

with  $\mathbf{h}(\xi)$  the solution to the ordinary differential equation

$$\begin{aligned} \partial_\xi \mathbf{h}(\xi) &= \mathbf{r}_k(\mathbf{h}(\xi)), & \xi \in \mathbb{R}, \\ \mathbf{h}(\xi_L) &= \mathbf{q}_L, \end{aligned} \quad (2.7)$$

for some  $\xi_L \in \mathbb{R}$ .

Furthermore, to each  $k$ -path corresponds a set of functions which are invariant on  $\mathcal{R}_k$ :

**Definition 2.5** Consider the hyperbolic system (2.1a). Let  $\mathbf{r}_k(\mathbf{q})$  denote the  $k^{\text{th}}$  eigenvector of the Jacobian  $\mathbf{A}(\mathbf{q}) = \partial_{\mathbf{q}} \mathbf{f}(\mathbf{q})$ . A  $k$ -Riemann invariant is any function  $\psi_k \in C^1(\mathbb{R}^n, \mathbb{R})$  satisfying

$$\partial_{\mathbf{q}} \psi_k(\mathbf{q}) \cdot \mathbf{r}_k(\mathbf{q}) = 0, \quad \forall \mathbf{q} \in \mathbb{R}^n. \quad (2.8)$$

There are at most  $n-1$  such  $k$ -Riemann invariants with linearly independent gradients in  $\mathbb{R}^n$ . Observe that for a linearly degenerate eigenpair  $(\lambda_k(\mathbf{q}), \mathbf{r}_k(\mathbf{q}))$  the eigenvalue  $\lambda_k(\mathbf{q})$  is a  $k$ -Riemann invariant.

## 2.2 Solution

The general solution to (2.1) consists of regions in the  $(x, t)$ -domain where the solution is constant, separated by simple waves, contact discontinuities and shock waves. Before constructing the general solution, we first obtain the (weak) solution to (2.1) in case that it contains only one of the aforementioned contact phenomena.

We establish that the (weak) solution to the Riemann problem can generally be written in similarity form (see, e.g., [Smo83]):

**Theorem 2.1** Suppose a unique solution  $\mathbf{q}(x, t)$  to the Riemann Problem (2.1) exists. Then  $\mathbf{q}(x, t)$  can be written in similarity form  $\mathbf{q}(x, t) = \mathbf{h}(x/t)$ .

**Proof:** Assume  $\mathbf{q}(x, t)$  solves (2.1). Then for all  $\alpha \in \mathbb{R}$ ,  $\mathbf{q}(\alpha x, \alpha t)$  is also a solution:

$$\partial_t \mathbf{q}(\alpha x, \alpha t) + \partial_x \mathbf{f}(\mathbf{q}(\alpha x, \alpha t)) = \alpha [\mathbf{D}_2 \mathbf{q}(\alpha x, \alpha t) + \mathbf{A}(\mathbf{q}(\alpha x, \alpha t)) \cdot \mathbf{D}_1 \mathbf{q}(\alpha x, \alpha t)] = 0, \quad (2.9)$$

where  $\mathbf{D}_l$  denotes differentiation with respect to the  $l^{\text{th}}$  function-argument. Because the solution is unique by assumption,  $\mathbf{q}(x, t) = \mathbf{q}(\alpha x, \alpha t)$ . Hence,  $\mathbf{q}(x, t) = \mathbf{h}(x/t)$ .  $\square$

A (classical) simple wave solution of (2.1) exists if  $\lambda_k(\mathbf{q})$  is a genuinely nonlinear eigenvalue,  $\lambda_k(\mathbf{q}_L) < \lambda_k(\mathbf{q}_R)$  and  $\mathbf{q}_R$  is on the  $k$ -path through  $\mathbf{q}_L$ . Note that this implies that the  $k$ -Riemann invariants are equal for  $\mathbf{q}_L$  and  $\mathbf{q}_R$ , i.e.,  $\psi_k^m(\mathbf{q}_L) = \psi_k^m(\mathbf{q}_R)$ , for  $m \neq k$ ,  $m = 1, \dots, n$ . Assuming that the genuinely nonlinear eigenvector in (2.7) is normalised such that

$$\partial_{\mathbf{q}} \lambda_k(\mathbf{q}) \cdot \mathbf{r}_k(\mathbf{q}) = 1, \quad \forall \mathbf{q} \in \mathbb{R}^n, \quad (2.10)$$

we find that  $\mathbf{q}(x, t) = \mathbf{h}(x/t)$  is the similarity solution in the simple wave region  $\lambda_k(\mathbf{q}_L) < x/t < \lambda_k(\mathbf{q}_R)$  (see, e.g., [Smo83], [Lax73]):

**Theorem 2.2** Suppose  $\mathbf{h} \in C^1(\mathbb{R}, \mathbb{R}^n)$  solves (2.7), with  $\mathbf{r}_k(\mathbf{q})$  normalised according to (2.10), and  $\mathbf{q}_R \in \mathcal{R}_k(\mathbf{q}_L)$ . Then  $\mathbf{q}(x, t) = \mathbf{h}(x/t)$  is the similarity solution of (2.1) in the simple-wave region  $\lambda_k(\mathbf{q}_L) < x/t < \lambda_k(\mathbf{q}_R)$ .

**Proof:** We will only show that  $\mathbf{q}(x, t) = \mathbf{h}(x/t)$  solves (2.1a). Inserting  $\mathbf{q}(x, t) = \mathbf{h}(x/t)$  in (2.1a), one obtains

$$\partial_t \mathbf{h}(x/t) + \partial_x \mathbf{f}(\mathbf{h}(x/t)) = \frac{1}{t} \left( \mathbf{A}(\mathbf{h}(x/t)) - \mathbf{I} \frac{x}{t} \right) \cdot \mathbf{Dh}(x/t), \quad (2.11)$$

where  $\mathbf{I}$  stands for the  $\mathbb{R}^{n \times n}$  identity matrix and  $\mathbf{D}$  denotes differentiation with respect to the function-argument. The right hand side term of (2.11) vanishes if  $x/t = \lambda_k(\mathbf{h}(x/t))$  and  $\mathbf{Dh}(x/t) = \mathbf{r}_k(\mathbf{h}(x/t))$ . The latter trivially follows from (2.7), the former from (2.10). Hence,  $\mathbf{h}(x/t)$  solves (2.1a).  $\square$

Outside the wave region the solution remains unchanged. The Riemann solution  $\mathbf{q}(x, t)$  in the case of a  $k$ -rarefaction wave is now simply composed of the solutions on the separate regions:

$$\mathbf{q}(x, t) = \begin{cases} \mathbf{q}_L, & \text{if } x/t < \lambda(\mathbf{q}_L), \\ \mathbf{h}(x/t), & \text{if } \lambda(\mathbf{q}_L) < x/t < \lambda(\mathbf{q}_R), \\ \mathbf{q}_R, & \text{if } x/t > \lambda(\mathbf{q}_R). \end{cases} \quad (2.12)$$

Next, we derive the (weak) Riemann solution in the instance that it contains a single contact discontinuity. The states  $\mathbf{q}_L$  and  $\mathbf{q}_R$  are connected by a  $k$ -contact discontinuity if  $(\lambda_k(\mathbf{q}), \mathbf{r}_k(\mathbf{q}))$  is a linearly degenerate eigenpair and  $\mathbf{q}_R$  is on the  $k$ -path through  $\mathbf{q}_L$ . Then, by (2.4),  $\lambda_k(\mathbf{q}_R) = \lambda_k(\mathbf{q}_L)$ . The solution to the Riemann problem is now obtained immediately from (2.12):

$$\mathbf{q}(x, t) = \begin{cases} \mathbf{q}_L, & \text{if } x/t < \lambda(\mathbf{q}_L) = \lambda(\mathbf{q}_R), \\ \mathbf{q}_R, & \text{if } x/t > \lambda(\mathbf{q}_L) = \lambda(\mathbf{q}_R). \end{cases} \quad (2.13)$$

However, because (2.13) is discontinuous at  $x/t = \lambda(\mathbf{q}_L) = \lambda(\mathbf{q}_R)$ , it must be verified that (2.13) satisfies the weak form of (2.1a):

$$\oint_{\mathcal{C}} \mathbf{q} n_t + \mathbf{f}(\mathbf{q}) n_x d\mathcal{C} = 0. \quad (2.14)$$

Here  $\mathcal{C}$  is any closed curve in  $(x, t)$  and  $\mathbf{n} = (n_t, n_x)$  denotes the outward pointing unit normal on  $\mathcal{C}$ . It can easily be shown that (2.14) does indeed hold for (2.13), so that (2.13) is a valid weak solution.

Finally, we consider the solution to (2.1) when it comprises a single shock. A shock occurs if  $\lambda_k(\mathbf{q})$  is a genuinely nonlinear eigenvalue,  $\lambda_k(\mathbf{q}_L) > \lambda_k(\mathbf{q}_R)$  and  $\mathbf{q}_R$  is on the  $k$ -shock path through  $\mathbf{q}_L$ . A solution of the form (2.12) is then necessarily multiple-valued and must therefore be discarded. Instead, the weak solution reads

$$\mathbf{q}(x, t) = \begin{cases} \mathbf{q}_L, & \text{if } x/t < s(\mathbf{q}_L; \mathbf{q}_R), \\ \mathbf{q}_R, & \text{if } x/t > s(\mathbf{q}_L; \mathbf{q}_R), \end{cases} \quad (2.15)$$

where  $s(\mathbf{q}_L; \mathbf{q}_R)$  denotes the shock speed, determined by the Rankine-Hugoniot relation

$$s(\mathbf{q}_L; \mathbf{q}_R)(\mathbf{q}_L - \mathbf{q}_R) = \mathbf{f}(\mathbf{q}_L) - \mathbf{f}(\mathbf{q}_R). \quad (2.16)$$

Expression (2.16) is in fact equivalent to (2.14). Hence, (2.15) is a valid weak solution of (2.1).

The general solution to the Riemann problem consists of  $n + 1$  (possibly empty) regions  $\Omega_l$  where the solution is constant, separated by simple waves, contact discontinuities and shock waves. Define  $\mathbf{q}_0 = \mathbf{q}_L$ ,  $\mathbf{q}_1 = \mathbf{q}_R$  and let  $\mathbf{q}_{l/n}$ ,  $l = 0, \dots, n$ , be the solution in  $\Omega_l$ . Assuming that  $\mathbf{q}_{(l-1)/n}$  is connected to  $\mathbf{q}_{l/n}$  by a simple wave, we denote by  $\mathbf{h}_l(x/t)$  the similarity solution in the wave region. Conversely, if  $\mathbf{q}_{(l-1)/n}$  is connected to  $\mathbf{q}_{l/n}$  by a shock wave, we designate  $s_l$  the appropriate shock speed. Then, in succinct form:

$$\mathbf{q}(x, t) = \begin{cases} \mathbf{q}_0, & \text{if } x/t < \sigma_0^+, \\ \mathbf{q}_{l/n}, & \text{if } \sigma_l^- < x/t < \sigma_l^+, \quad l = 1, \dots, n-1, \\ \mathbf{h}_l(x/t), & \text{if } \sigma_{l-1}^+ < x/t < \sigma_l^-, \quad l = 1, \dots, n-1, \\ \mathbf{q}_1, & \text{if } x/t > \sigma_n^-, \end{cases} \quad (2.17a)$$

where  $\sigma_l^\pm$  denotes the contact speed

$$\sigma_l^\pm = \begin{cases} \lambda_{l+(1\pm 1)/2}(\mathbf{q}_{l/n}) & \text{if } \pm \lambda_{l+(1\pm 1)/2}(\mathbf{q}_{l/n}) < \pm \lambda_{l+(1\pm 1)/2}(\mathbf{q}_{(l\pm 1)/n}), \\ s_{l+(1\pm 1)/2} & \text{otherwise.} \end{cases} \quad (2.17b)$$

The general solution (2.17) is schematically depicted in figure 1. The figure illustrates the contiguity of regions connected by shock waves and contact discontinuities, for instance,  $\Omega_{(l-1)/n}$  and  $\Omega_{l/n}$ , and the separation of regions connected by rarefaction waves, e.g.,  $\Omega_{l/n}$  and  $\Omega_{(l+1)/n}$ .

As a sidenote, we mention that for general  $\mathbf{f}(\mathbf{q})$  and sufficiently large  $\|\mathbf{q}_L - \mathbf{q}_R\|$ , a solution to (2.1) can be non-existent (see, for example, [Smo83]).




Figure 1: General solution to the Riemann problem

### 3. APPROXIMATE RIEMANN SOLUTION

In the previous section we established that the solution to the Riemann problem can generally be written in similarity form  $\mathbf{h}(x/t)$ . Denoting by  $\mathbf{h}(x/t; \mathbf{q}_L, \mathbf{q}_R)$  the similarity solution for given  $\mathbf{q}_L$  and  $\mathbf{q}_R$ , we find  $\mathbf{f}(\mathbf{h}(0; \mathbf{q}_L, \mathbf{q}_R))$  to be the corresponding centered flux,  $\mathbf{f}(\mathbf{q}_L, \mathbf{q}_R)$ . This flux is of particular importance in computational applications: following Godunov's approach, it can be interpreted as the flux between two adjacent cells in the discretised domain. Unfortunately, solving the Riemann problem exactly is computationally expensive and it is therefore necessary to revert to approximate solution techniques.

In this section, we investigate Osher's approximate Riemann solver and a modified Osher-type scheme. We will first present a general outline of the Osher scheme. Subsequently, the approximate Riemann solution employed in Osher's scheme is examined and the computed flux approximation is

compared to the exact solution. Finally, we shall propose the modified scheme, based on the preceding analysis.

### 3.1 Osher's scheme

In the scheme developed by Osher [OS82], [OC83], the centered flux  $\mathbf{f}(\mathbf{q}_L, \mathbf{q}_R) = \mathbf{f}(\mathbf{h}(0; \mathbf{q}_L, \mathbf{q}_R))$  is approximated by:

$$\tilde{\mathbf{f}}(\mathbf{q}_L, \mathbf{q}_R) = \frac{1}{2}\mathbf{f}(\mathbf{q}_L) + \frac{1}{2}\mathbf{f}(\mathbf{q}_R) - \frac{1}{2} \int_{\mathbf{q}_L}^{\mathbf{q}_R} |\mathbf{A}(\mathbf{w})| \cdot d\mathbf{w}, \quad (3.1)$$

with the absolute value of the Jacobian matrix  $\mathbf{A}(\mathbf{q})$  defined by  $|\mathbf{A}(\mathbf{q})| \equiv \mathbf{R}(\mathbf{q}) \cdot |\boldsymbol{\Lambda}(\mathbf{q})| \cdot \mathbf{R}(\mathbf{q})^{-1}$ . Here,  $|\boldsymbol{\Lambda}(\mathbf{q})| = \text{diag}(|\lambda_1(\mathbf{q})|, \dots, |\lambda_n(\mathbf{q})|)$ . Clearly, the integral term represents the upwind contribution to the centered flux approximation.

The integral in (3.1) is evaluated along a path  $\Gamma = \{\mathbf{q}(s) : 0 \leq s \leq 1\} \subset \mathbb{R}^n$  in state space, satisfying  $\mathbf{q}(0) = \tilde{\mathbf{q}}_0$  and  $\mathbf{q}(1) = \tilde{\mathbf{q}}_1$ , with  $\tilde{\mathbf{q}}_0 = \mathbf{q}_L$  and  $\tilde{\mathbf{q}}_1 = \mathbf{q}_R$  or vice versa. This path is composed of sub-paths  $\Gamma_l$ ,  $l = 1, 2, \dots, n$ , where each of the sub-paths connects two adjacent states  $\tilde{\mathbf{q}}_{(l-1)/n}$  and  $\tilde{\mathbf{q}}_{l/n}$ . Moreover,  $\Gamma_l$  is tangential to an eigenvector  $\mathbf{r}_{k(l)}$ , where  $k : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  is a bijective mapping. It should be appreciated here that  $\Gamma_l$  is thus a section of the  $k(l)$ -path through  $\tilde{\mathbf{q}}_{(l-1)/n}$ , connecting  $\tilde{\mathbf{q}}_{(l-1)/n}$  and  $\tilde{\mathbf{q}}_{l/n}$ . Usual choices for the ordering of the sub-paths are the O-variant  $k(l) = n - l$  and the P-variant  $k(l) = l$ .

The integral term in (3.1) can be rewritten as a summation of the contributions of the integral over each of the sub-paths:

$$\int_{\mathbf{q}_L}^{\mathbf{q}_R} |\mathbf{A}(\mathbf{w})| \cdot d\mathbf{w} = \sum_{l=1}^n \int_{\Gamma_l} |\mathbf{A}(\mathbf{w}(\xi))| \cdot \mathbf{r}_{k(l)}(\mathbf{w}(\xi)) d\xi = \sum_{l=1}^n \int_{\Gamma_l} \text{sign}(\lambda_{k(l)}(\mathbf{w})) \mathbf{A}(\mathbf{w}) \cdot d\mathbf{w} \quad (3.2)$$

Obviously, if  $\lambda_{k(l)}$  does not change sign along  $\Gamma_l$ , then the sub-integral can be evaluated to  $[\mathbf{f}(\tilde{\mathbf{q}}_{l/n}) - \mathbf{f}(\tilde{\mathbf{q}}_{(l-1)/n})] \text{sign}(\lambda_{k(l)})$ . Then, by (2.4), if  $\lambda_{k(l)} = \lambda_{k(l+1)} = \dots = \lambda_{k(l+\mu)}$  is a linearly degenerate eigenvalue, the sum in (3.2) concatenates and we simply obtain

$$\sum_{i=0}^{\mu} \int_{\Gamma_{l+i}} |\mathbf{A}(\mathbf{w})| \cdot d\mathbf{w} = \text{sign}(\lambda_{k(l)}(\mathbf{q}_{l/n})) [\mathbf{f}(\mathbf{q}_{l/n}) - \mathbf{f}(\mathbf{q}_{(l+\mu)/n})] \quad (3.3)$$

Hence, the intermediate stages  $\tilde{\mathbf{q}}_{(l+i)/n}$ ,  $i = 1, 2, \dots, \mu-1$  are of no consequence and may be eliminated from the composed path  $\Gamma$ .

As a result of the choice of the sub-paths  $\Gamma_l$ , the intermediate  $\tilde{\mathbf{q}}_{l/n}$ ,  $l = 1, 2, \dots, n-1$  can be conveniently determined by means of the Riemann invariants: Because the sub-path  $\Gamma_l \subset \mathcal{R}_{k(l)}(\tilde{\mathbf{q}}_{(l-1)/n})$ ,

$$\psi_{k(l)}^m(\tilde{\mathbf{q}}_{(l-1)/n}) = \psi_{k(l)}^m(\tilde{\mathbf{q}}_{l/n}), \quad l, m = 1, 2, \dots, n, \quad m \neq k(l), \quad (3.4)$$

see section 2.1. If it is assumed that the  $k$ -Riemann invariants in (3.4) have linearly independent gradients, then by the implicit function theorem, (3.4) constitutes a solvable system of equations from which the  $\tilde{\mathbf{q}}_{l/n}$ ,  $l = 1, 2, \dots, n$  can be extracted. In many practical cases the intermediate stages can then be solved explicitly from (3.4). Once the intermediate states  $\tilde{\mathbf{q}}_{l/n}$  have been obtained, the flux approximation  $\mathbf{f}(\mathbf{q}_L, \mathbf{q}_R)$  can be computed using (3.1), (3.2).



### 3.2 Accuracy

The flux computed by means of the Osher scheme,  $\tilde{\mathbf{f}}(\mathbf{q}_L, \mathbf{q}_R)$ , relies on an approximate solution of the Riemann problem. Because the approximation can again be written in similarity form, it is useful to introduce the notation  $\tilde{\mathbf{f}}(\mathbf{q}_L, \mathbf{q}_R) = \mathbf{f}(\tilde{\mathbf{h}}(0; \mathbf{q}_L, \mathbf{q}_R))$ , where  $\tilde{\mathbf{h}}(x/t; \mathbf{q}_L, \mathbf{q}_R)$  stands for the approximate similarity solution. In this section we investigate the accuracy of the approximate similarity solution and of the corresponding centered flux approximation.

To evaluate the accuracy of the approximate solution, we examine the inherent representation of simple waves, contact discontinuities and shock waves. In section 3.1 it was emphasized that the sub-paths,  $\Gamma_l$ , in Osher's scheme are actually sections of  $k(l)$ -paths. Referring to section 2.2, it follows that the intermediate states  $\tilde{\mathbf{q}}_{l/n}$ ,  $l = 0, \dots, n$ , in the approximate solution are connected by simple-waves only. Clearly, this representation is correct for simple waves and contact discontinuities. However, shock waves in the actual solution are then replaced by so-called *overtaken simple waves*, see [Lee84]. We will now show that this representation is accurate for weak shocks. From [Smo83] we adopt:

**Lemma 3.1** *Suppose  $\mathbf{q}_L$  and  $\mathbf{q}_R$  are connected by a weak  $k$ -shock with shock strength  $\epsilon$ , i.e.,  $\mathbf{q}_R \in \mathcal{S}_k(\mathbf{q}_L)$  and  $\lambda_k(\mathbf{q}_L) = \lambda_k(\mathbf{q}_R) + \epsilon$ , with  $\epsilon$  a small positive number. Then the change in a  $k$ -Riemann invariant across the  $k$ -shock is of order  $\mathcal{O}(\epsilon^3)$ .*

**Proof:** *Proof is omitted here, but can be found in [Smo83, pages 326–333].*  $\square$

Then, we obtain:

**Theorem 3.1** *Suppose  $\mathbf{q}_R \in \mathcal{S}_k(\mathbf{q}_L)$  and  $\lambda_k(\mathbf{q}_L) = \lambda_k(\mathbf{q}_R) + \epsilon$ . Then a  $\tilde{\mathbf{q}}_R \in \mathcal{R}_k(\mathbf{q}_L)$  exists such that  $\lambda_k(\tilde{\mathbf{q}}_R) = \lambda_k(\mathbf{q}_R)$  and  $|\tilde{\mathbf{q}}_R - \mathbf{q}_R|$  is of order  $\mathcal{O}(\epsilon^3)$ .*

**Proof:** *By definition 2.5,  $\psi_k^m(\mathbf{q}_L) = \psi_k^m(\tilde{\mathbf{q}}_R)$ ,  $k = 1, 2, \dots, n$ ,  $k \neq m$ . Then, by lemma 3.1,*

$$\psi_k^m(\tilde{\mathbf{q}}_R) = \psi_k^m(\mathbf{q}_R) + \mathcal{O}(\epsilon^3). \quad (3.5)$$

*System (3.5) can be augmented with  $\lambda_k(\tilde{\mathbf{q}}_R) = \lambda_k(\mathbf{q}_R)$  to obtain  $n$  equations for  $\tilde{\mathbf{q}}_R$ . Because  $\text{rank}(\partial_{\mathbf{q}}\psi_k^1, \dots, \partial_{\mathbf{q}}\psi_k^n) = n - 1$  and  $\partial_{\mathbf{q}}\lambda_k \in (\partial_{\mathbf{q}}\psi_k^1, \dots, \partial_{\mathbf{q}}\psi_k^n)^\perp$ ,  $\det(\partial_{\mathbf{q}}\psi_k^1, \dots, \partial_{\mathbf{q}}\psi_k^n, \partial_{\mathbf{q}}\lambda_k) \neq 0$ . The result now simply follows by a Taylor expansion around  $\mathbf{q}_R$  of the terms in  $\tilde{\mathbf{q}}_R$  of the augmented system.*  $\square$

From theorem 3.1 it may be inferred that the intermediate states obtained by a rarefaction-waves-only approximation are  $\mathcal{O}(\epsilon_{\max}^3)$  accurate, with

$$\epsilon_{\max} = \max_{l=1 \dots n} (\lambda_l(\mathbf{q}_{(l-1)/n}) - \lambda_l(\mathbf{q}_{l/n}), 0) \quad (3.6)$$

the strength of the strongest shock.

Although the computed intermediate states are accurate even in the presence of (weak) shocks, the flux approximation  $\tilde{\mathbf{f}}(\mathbf{q}_L, \mathbf{q}_R)$  is not necessarily so. By (3.2), if  $\tilde{\mathbf{q}}_R \in \mathcal{R}_k(\mathbf{q}_L)$  and  $\lambda_k(\mathbf{q}_L) > 0 > \lambda_k(\tilde{\mathbf{q}}_R)$ ,

$$\tilde{\mathbf{f}}(\mathbf{q}_L, \tilde{\mathbf{q}}_R) = \mathbf{f}(\mathbf{q}_L) + \mathbf{f}(\tilde{\mathbf{q}}_R) - \mathbf{f}(\mathbf{q}^*), \quad (3.7)$$

with  $\mathbf{q}^* \in \mathcal{R}_k(\mathbf{q}_L)$  such that  $\lambda_k(\mathbf{q}^*) = 0$ . In contrast, ignoring terms  $\mathcal{O}(\epsilon^3)$ , one finds that the actual flux corresponding to the  $k$ -shock is  $\mathbf{f}(\mathbf{q}_L)$  if  $s(\mathbf{q}_R; \mathbf{q}_L) > 0$  and  $\mathbf{f}(\mathbf{q}_R)$  if  $s(\mathbf{q}_R; \mathbf{q}_L) < 0$ . Consequently, the error in the approximate flux in the instance of a centered shock with strength  $\epsilon$  may be of  $\mathcal{O}(\epsilon)$ .

### 3.3 Modified Scheme

In view of the above, a modification of the scheme is advocated. The rarefaction waves only approximation of the similarity solution is maintained. However, the centered flux approximation is obtained differently, to avoid loss of accuracy due to centered shock waves.

We propose to extract the intermediate states in the approximate solution to the Riemann problem from

$$\psi_l^m(\tilde{\mathbf{q}}_{(l-1)/n}) = \psi_l^m(\tilde{\mathbf{q}}_{l/n}), \quad l, m = 1, 2, \dots, n, \quad m \neq l, \quad (3.8)$$

with  $\tilde{\mathbf{q}}_0 = \mathbf{q}_L$  and  $\tilde{\mathbf{q}}_1 = \mathbf{q}_R$ . This is in fact equivalent to (3.4) with a presumed P-variant ordering of the sub-paths. Next, approximate contact speeds  $\tilde{\sigma}_l^\pm$  are obtained:

$$\tilde{\sigma}_l^\pm = \begin{cases} \lambda_{l+(1\pm 1)/2}(\tilde{\mathbf{q}}_{l/n}) & \text{if } \pm \lambda_{l+(1\pm 1)/2}(\tilde{\mathbf{q}}_{l/n}) < \pm \lambda_{l+(1\pm 1)/2}(\tilde{\mathbf{q}}_{(l\pm 1)/n}), \\ \tilde{s}_{l+(1\pm 1)/2} & \text{otherwise,} \end{cases} \quad (3.9a)$$

with

$$\tilde{s}_{l+(1\pm 1)/2} = \frac{1}{2}\lambda_{l+(1\pm 1)/2}(\tilde{\mathbf{q}}_{l/n}) + \frac{1}{2}\lambda_{l+(1\pm 1)/2}(\tilde{\mathbf{q}}_{(l\pm 1)/n}). \quad (3.9b)$$

Estimate (3.9b) of the shock speed is justified by the following theorem, taken from [Smo83]:

**Theorem 3.2** *Suppose  $\mathbf{q}_R \in \mathcal{S}_k(\mathbf{q}_L)$  and  $\lambda_k(\mathbf{q}_L) = \lambda_k(\mathbf{q}_R) + \epsilon$ ,  $\epsilon > 0$ . Then the speed of the  $k$ -shock connecting  $\mathbf{q}_L$  and  $\mathbf{q}_R$  satisfies  $s(\mathbf{q}_L; \mathbf{q}_R) = \frac{1}{2}\lambda_k(\mathbf{q}_L) + \frac{1}{2}\lambda_k(\mathbf{q}_R) + \mathcal{O}(\epsilon^2)$ .*

**Proof:** *Proof can be found in [Smo83, pages 326–333].*  $\square$

Once the intermediate states and contact speeds have been established, the approximate Riemann solution can be constructed in a manner similar to (2.17a). However, considering that our purpose is to compute an approximation to the centered flux, we only need to obtain the central part of the approximate solution:

$$\tilde{\mathbf{h}}(0; \mathbf{q}_L, \mathbf{q}_R) = \begin{cases} \tilde{\mathbf{q}}_0, & \text{if } \tilde{\sigma}_0^+ > 0, \\ \tilde{\mathbf{q}}_{l/n}, & \text{if } \tilde{\sigma}_l^- < 0 < \tilde{\sigma}_l^+, \quad l \in \{1, \dots, n-1\}, \\ \tilde{\mathbf{q}}^*, & \text{if } \tilde{\sigma}_{l-1}^+ < 0 < \tilde{\sigma}_l^-, \quad l \in \{1, \dots, n-1\}, \\ \tilde{\mathbf{q}}_1, & \text{if } \tilde{\sigma}_n^- < 0, \end{cases} \quad (3.10)$$

with  $\tilde{\mathbf{q}}^* \in \mathcal{R}_l(\tilde{\mathbf{q}}_{(l-1)/n})$  such that  $\lambda_l(\tilde{\mathbf{q}}^*) = 0$  in case of a centered rarefaction wave. The centered flux approximation is now simply  $\tilde{\mathbf{f}}(\mathbf{q}_L, \mathbf{q}_R) = \mathbf{f}(\tilde{\mathbf{h}}(0; \mathbf{q}_L, \mathbf{q}_R))$ .

#### 4. APPLICATIONS IN HYDRODYNAMICS

In the previous section we presented a flux-difference splitting scheme that gives an accurate approximation of the centered flux in the Riemann problem, even in the presence of (weak) centered shock waves. A prerequisite for the flux evaluation is the derivation of the intermediate states  $\tilde{\mathbf{q}}_{l/n}$ ,  $l = 1, \dots, n$ . Once these states have been obtained, the flux calculation proceeds via straightforward operations.

In this section we derive the intermediate states for the one dimensional Euler equations for three types of fluids that are used to model the behaviour of water. These fluids are, successively, a genuinely compressible fluid, an artificially compressible fluid and an incompressible fluid. Furthermore, we obtain the intermediate states for the Euler equations in the case of an immiscible, compressible two-phase flow.

##### 4.1 Compressible fluid

Suppose that  $u$ ,  $v$  and  $w$  denote the  $x$ ,  $y$  and  $z$  components of a fluid velocity  $\mathbf{u} \in \mathbb{R}^3$  in a Cartesian coordinate system, respectively, and that  $\rho \in \mathbb{R}^+$  denotes the density of the fluid. Consider the hyperbolic system (2.1a) with  $\mathbf{q} = (\rho u, \rho v, \rho w, \rho)^T$  and  $\mathbf{f}(\mathbf{q})$  given by

$$\mathbf{f}(\mathbf{q}) = (q_1^2/q_4 + p(q_4), q_1 q_2/q_4, q_1 q_3/q_4, q_1)^T. \quad (4.1)$$

Then equations (2.1a) are the Euler equations for a compressible fluid in one dimension. In this section it is assumed that the pressure is related to the density via an equation of state of the form  $p = p(\rho)$ , with  $p \in C^1(\mathbb{R}^+, \mathbb{R}^+)$  an increasing function. An example is Tait's equation of state, which is often used to model the behaviour of water:

$$p(\rho) = \alpha\rho^\gamma + \beta, \quad (4.2)$$

where  $\alpha, \gamma \in ]0, \infty[$  and  $\beta \in \mathbb{R}$  are given constants. Our objective now is to obtain the approximate intermediate states for the Euler equations (2.1a), (4.1).

In order to compute the intermediate states from (3.8),  $k$ -Riemann invariants for the system under consideration have to be derived first. The Jacobian of the flux vector (4.1) reads

$$\mathbf{A}(\mathbf{q}) = \partial_{\mathbf{q}}\mathbf{f} = \begin{pmatrix} 2q_1/q_4 & 0 & 0 & -q_1^2/q_4^2 + c^2(q_4) \\ q_2/q_4 & q_1/q_4 & 0 & -q_1q_2/q_4 \\ q_3/q_4 & 0 & q_1/q_4 & -q_3q_1/q_4 \\ 1 & 0 & 0 & 0 \end{pmatrix}, \quad (4.3)$$

where  $c(\rho) = \sqrt{\partial_\rho p}$  denotes the speed of sound. Computation of the eigenvalues of  $\mathbf{A}(\mathbf{q})$  and the corresponding eigenvectors then yields

$$\lambda_1 = q_1/q_4 - c(q_4), \lambda_{2,3} = q_1/q_4, \lambda_4 = q_1/q_4 + c(q_4), \quad (4.4)$$

and

$$\begin{aligned} \mathbf{r}_1 &= (q_1/q_4 - c(q_4), q_2/q_4, q_3/q_4, 1)^T, \\ \mathbf{r}_2 &= (0, 1, 0, 0)^T, \\ \mathbf{r}_3 &= (0, 0, 1, 0)^T, \\ \mathbf{r}_4 &= (q_1/q_4 + c(q_4), q_2/q_4, q_3/q_4, 1)^T. \end{aligned} \quad (4.5)$$

Notice that the eigenvalue  $\lambda_k$  and the eigenvector  $\mathbf{r}_k$  are genuinely nonlinear for  $k = 1, 4$  and linearly degenerate for  $k = 2, 3$ . Riemann invariants are then obtained by solving partial differential equations (2.8), with the eigenvectors according to (4.5). The details are omitted here, but it can easily be verified that (4.6) constitutes a complete set of  $k$ -Riemann invariants:

$$\begin{aligned} \psi_1^2 &= q_1/q_4 + \phi(q_4), & \psi_1^3 &= q_2/q_4, & \psi_1^4 &= q_3/q_4, \\ \psi_2^1 &= q_1, & \psi_2^3 &= q_3, & \psi_2^4 &= q_4, \\ \psi_3^1 &= q_1, & \psi_3^2 &= q_2, & \psi_3^4 &= q_4, \\ \psi_4^1 &= q_1/q_4 - \phi(q_4), & \psi_4^2 &= q_2/q_4, & \psi_4^3 &= q_3/q_4, \end{aligned} \quad (4.6a)$$

where  $\phi(\rho)$  is defined

$$\phi(\rho) = \int_0^\rho \frac{c(\eta)}{\eta} d\eta. \quad (4.6b)$$

The intermediate states can now be extracted from (3.8), (4.6). In view of the linear degeneracy of the eigenvalues  $\lambda_2$  and  $\lambda_3$  and the arguments presented in section 3.1, we ignore  $\tilde{\mathbf{q}}_{1/2}$ . We then find that  $\mathbf{q}_0$  is connected to  $\mathbf{q}_1$  via two approximate intermediate states  $\tilde{\mathbf{q}}_{1/3}$  and  $\tilde{\mathbf{q}}_{2/3}$ :

$$\tilde{\mathbf{q}}_{1/3} = \phi^{-1} \left( \frac{1}{2}\psi_1^2(\mathbf{q}_0) - \frac{1}{2}\psi_4^1(\mathbf{q}_1) \right) \begin{pmatrix} \frac{1}{2}\psi_1^2(\mathbf{q}_0) + \frac{1}{2}\psi_4^1(\mathbf{q}_1) \\ \psi_1^3(\mathbf{q}_0) \\ \psi_1^4(\mathbf{q}_0) \\ 1 \end{pmatrix} \quad (4.7)$$

and

$$\tilde{\mathbf{q}}_{2/3} = \phi^{-1} \left( \frac{1}{2}\psi_1^2(\mathbf{q}_0) - \frac{1}{2}\psi_4^1(\mathbf{q}_1) \right) \begin{pmatrix} \frac{1}{2}\psi_1^2(\mathbf{q}_0) + \frac{1}{2}\psi_4^1(\mathbf{q}_1) \\ \psi_4^2(\mathbf{q}_1) \\ \psi_4^3(\mathbf{q}_1) \\ 1 \end{pmatrix}, \quad (4.8)$$

where  $\phi^{-1}(\psi)$  denotes the inverse of  $\phi(\psi)$ .

For a fluid that is described by Tait's equation of state, the intermediate states can be determined by substituting (4.2) in equations (4.6) to (4.8). The intermediate velocity components  $\tilde{v}_{1/3}$ ,  $\tilde{v}_{2/3}$ ,  $\tilde{w}_{1/3}$  and  $\tilde{w}_{2/3}$  are immediately obtained from (4.6):

$$\begin{aligned} \tilde{v}_{1/3} &= v_0, & \tilde{v}_{2/3} &= v_1, \\ \tilde{w}_{1/3} &= w_0, & \tilde{w}_{2/3} &= w_1. \end{aligned} \quad (4.9)$$

From (4.6) it is also clear that  $\tilde{u}_{1/3} = \tilde{u}_{2/3} \equiv \tilde{u}_{1/2}$  and  $\tilde{\rho}_{1/3} = \tilde{\rho}_{2/3} \equiv \tilde{\rho}_{1/2}$ . To determine  $\tilde{u}_{1/2}$  and  $\tilde{\rho}_{1/2}$ , it is necessary to distinguish between the cases  $\gamma = 1$  and  $\gamma \neq 1$ . For  $\gamma = 1$  one obtains

$$\begin{aligned} \tilde{u}_{1/2} &= \frac{1}{2}(u_0 + u_1) + \frac{\alpha}{2} \ln(\rho_0/\rho_1), \\ \tilde{\rho}_{1/2} &= \sqrt{\rho_0\rho_1} \exp\left(\frac{u_0 + u_1}{2\alpha}\right). \end{aligned} \quad (4.10)$$

In case  $\gamma \neq 1$ , it is convenient to express the density in terms of the speed of sound:

$$\begin{aligned} \tilde{u}_{1/2} &= \frac{1}{2}(u_0 + u_1) + \frac{1}{\gamma - 1}[c(\rho_0) - c(\rho_1)], \\ c(\tilde{\rho}_{1/2}) &= \frac{\gamma - 1}{4}(u_0 - u_1) + \frac{1}{2}[c(\rho_0) + c(\rho_1)]. \end{aligned} \quad (4.11)$$

#### 4.2 Artificially compressible fluid

Assume that  $u$ ,  $v$  and  $w$  again denote the  $x$ ,  $y$  and  $z$  components of a fluid velocity  $\mathbf{u} \in \mathbb{R}^3$  in a Cartesian coordinate system, respectively, and that  $p \in \mathbb{R}^+$  denotes the fluid pressure. Consider hyperbolic system (2.1a) with  $\mathbf{q} = (u, v, w, p)^T$ . Let  $\mathbf{f}(\mathbf{q})$  be

$$\mathbf{f}(\mathbf{q}) = (q_1^2 + q_4, q_1q_2, q_1q_3, c^2q_1)^T, \quad (4.12)$$

with  $c$  constant. Equations (2.1a), (4.12) are the Euler equations for an artificially compressible fluid in one dimension. Notice that the  $\partial_t p$  term that occurs in (2.1a) in this case, implies compressibility of the fluid.

To obtain the intermediate states, we first derive Riemann invariants for (2.1a), (4.12). For the Jacobian of  $\mathbf{f}(\mathbf{q})$  we simply obtain

$$\mathbf{A}(\mathbf{q}) = \partial_{\mathbf{q}}\mathbf{f} = \begin{pmatrix} 2q_1 & 0 & 0 & 1 \\ q_2 & q_1 & 0 & 0 \\ q_3 & 0 & q_1 & 0 \\ c^2 & 0 & 0 & 0 \end{pmatrix}. \quad (4.13)$$

The eigenvalues of  $\mathbf{A}(\mathbf{q})$  and the corresponding eigenvectors follow by straightforward computation:

$$\lambda_1 = q_1 - \sqrt{q_1^2 + c^2}, \lambda_{2,3} = q_1, \lambda_4 = q_1 + \sqrt{q_1^2 + c^2}, \quad (4.14)$$

and

$$\begin{aligned}
\mathbf{r}_1 &= \left(1, -q_2/\sqrt{q_1^2 + c^2}, -q_3/\sqrt{q_1^2 + c^2}, -q_1 - \sqrt{q_1^2 + c^2}\right)^T, \\
\mathbf{r}_2 &= (0, 1, 0, 0)^T, \\
\mathbf{r}_3 &= (0, 0, 1, 0)^T, \\
\mathbf{r}_4 &= \left(1, q_2/\sqrt{q_1^2 + c^2}, q_3/\sqrt{q_1^2 + c^2}, -q_1 + \sqrt{q_1^2 + c^2}\right)^T.
\end{aligned} \tag{4.15}$$

The eigenpairs  $(\lambda_1, \mathbf{r}_1)$  and  $(\lambda_4, \mathbf{r}_4)$  are genuinely non-linear, whereas the eigenpairs  $(\lambda_2, \mathbf{r}_2)$  and  $(\lambda_3, \mathbf{r}_3)$  are linearly degenerate. Riemann invariants are now obtained by solving (2.8), (4.15):

$$\begin{aligned}
\psi_1^2 &= q_2 \lambda_4, & \psi_1^3 &= q_3 \lambda_4, & \psi_1^4 &= \lambda_4 \exp([2q_4 + q_1 \lambda_4]/c^2), \\
\psi_2^1 &= q_1, & \psi_2^3 &= q_3, & \psi_2^4 &= q_4, \\
\psi_3^1 &= q_1, & \psi_3^2 &= q_2, & \psi_3^4 &= q_4, \\
\psi_4^1 &= q_2 \lambda_1, & \psi_4^2 &= q_3 \lambda_1, & \psi_4^3 &= \lambda_1 \exp([2q_4 + q_1 \lambda_1]/c^2).
\end{aligned} \tag{4.16}$$

The foregoing invariants have linearly independent gradients. Hence, the intermediate states can be obtained from (3.8), (4.16).

Considering the linear degeneracy of  $\lambda_2, \lambda_3$ , we only need to obtain  $\tilde{\mathbf{q}}_{1/3}$  and  $\tilde{\mathbf{q}}_{2/3}$ . Unfortunately, in this instance we have not succeeded in deriving a closed form expression for these intermediate states. However, from (4.16) it immediately follows that  $\tilde{u}_{1/3} = \tilde{u}_{2/3} \equiv \tilde{u}_{1/2}$  and  $\tilde{\rho}_{1/3} = \tilde{\rho}_{2/3} \equiv \tilde{\rho}_{1/2}$ . Then, using the expressions for  $\psi_1^4$  and  $\psi_4^3$ , one finds that  $\tilde{u}_{1/2}$  is determined by the implicit relation:

$$\left( \frac{\tilde{u}_{1/2} + \sqrt{\tilde{u}_{1/2}^2 + c^2}}{\tilde{u}_{1/2} - \sqrt{\tilde{u}_{1/2}^2 + c^2}} \right) \exp\left( \frac{2\tilde{u}_{1/2}\sqrt{\tilde{u}_{1/2}^2 + c^2}}{c^2} \right) = \frac{\psi_1^4(\mathbf{q}_0)}{\psi_4^3(\mathbf{q}_1)}, \tag{4.17}$$

Once  $\tilde{u}_{1/2}$  has been solved from (4.17),  $\tilde{\mathbf{q}}_{1/3}$  and  $\tilde{\mathbf{q}}_{2/3}$  are simply obtained from (4.16).

### 4.3 Incompressible fluid

We commence by analysing the Euler equations for an incompressible flow. Assume that  $\mathbf{u} \in \mathbb{R}^3$  denotes the fluid velocity and that the fluid pressure divided by the (constant) fluid density is designated  $p \in \mathbb{R}^+$ . Next, let  $\mathbf{F} \in C^1(\mathbb{R}^3, \mathbb{R}^{3 \times 3})$  be the convective momentum flux tensor, defined by the dyad  $\mathbf{F}(\mathbf{u}) = \mathbf{u}\mathbf{u}$ . The Euler equations for an incompressible fluid read

$$\partial_t \mathbf{u} + \mathbf{F}(\mathbf{u}) \cdot \tilde{\nabla} + \tilde{\nabla} p = 0, \tag{4.18a}$$

$$\tilde{\nabla} \cdot \mathbf{u} = 0, \tag{4.18b}$$

where the postgradient operator is defined  $\mathbf{u}\tilde{\nabla} = (\tilde{\nabla}\mathbf{u})^T$ . Due to the absence of a time derivative in (4.18b), equations (4.18) do not constitute a hyperbolic system. However, equations (4.18) can be rewritten as an inhomogeneous hyperbolic system governing  $\mathbf{u}$  and a Poisson equation for  $p$ . Indeed, inserting  $\mathbf{F}(\mathbf{u})$  and replacing (4.18b) by the divergence of (4.18a) with (4.18b) implicitly imposed, we obtain:

$$\partial_t \mathbf{u} + \mathbf{u}\mathbf{u} \cdot \tilde{\nabla} = -\tilde{\nabla} p, \tag{4.19a}$$

$$\Delta p = -\mathbf{u}\tilde{\nabla} : \tilde{\nabla}\mathbf{u}, \tag{4.19b}$$

where  $\Delta$  denotes the Laplace operator in  $\mathbb{R}^3$ . Solving the Euler equations for an incompressible fluid now requires the simultaneous resolution of the hyperbolic system (4.19a) and the elliptic Poisson equation (4.19b). Here, we shall only concern ourselves with the hyperbolic part of the operator. Furthermore, in the following section we will only consider the homogeneous system in one dimension,

i.e., we shall neglect the source term  $-\vec{\nabla}p$  and (assuming a Cartesian coordinate system is employed) the flux gradients in the  $y$  and  $z$  direction. We then retrieve an expression of the form (2.1a), with  $\mathbf{q} = (u, v, w)^T$ , where  $u, v, w$  again denote the  $x, y, z$  components of the fluid velocity  $\mathbf{u} \in \mathbb{R}^3$  in a Cartesian coordinate system, respectively, and  $\mathbf{f}(\mathbf{q})$  given by

$$\mathbf{f}(\mathbf{q}) = (q_1^2, q_1 q_2, q_1 q_3)^T. \quad (4.20)$$

We appreciate that the first equation of (2.1a), (4.20) is decoupled from the remaining system and can therefore be treated separately. However, for completeness we refrain from doing so.

To obtain the approximate intermediate states for (2.1a), (4.20), we first determine Riemann invariants for this system. The Jacobian of  $\mathbf{f}(\mathbf{q})$  reads

$$\mathbf{A}(\mathbf{q}) = \partial_{\mathbf{q}} \mathbf{f} = \begin{pmatrix} 2q_1 & 0 & 0 \\ q_2 & q_1 & 0 \\ q_3 & 0 & q_1 \end{pmatrix}, \quad (4.21)$$

with the eigenvalues

$$\lambda_1 = q_1^2, \lambda_{2,3} = q_1, \quad (4.22)$$

and the corresponding eigenvectors

$$\begin{aligned} \mathbf{r}_1 &= (q_1, q_2, q_3)^T, \\ \mathbf{r}_2 &= (0, 1, 0)^T, \\ \mathbf{r}_3 &= (0, 0, 1)^T. \end{aligned} \quad (4.23)$$

The first eigenpair is neither linearly degenerate nor genuinely nonlinear: the gradient of  $\lambda_1(\mathbf{q})$  in the direction of  $\mathbf{r}_1(\mathbf{q})$  vanishes for  $q_1 = 0$ , but is nonzero otherwise. Nevertheless, for our purposes it is sufficient to treat  $(\lambda_1, \mathbf{r}_1)$  as a genuinely nonlinear eigenpair, because the eigenvalue vanishes only if  $q_1 = 0$  and, therefore, the eigenvalue can change sign only once along  $\mathcal{R}_1(\mathbf{q}_L)$ . The second and third eigenpair are linearly degenerate. Riemann invariants are obtained by solving (2.8), (4.23):

$$\begin{aligned} \psi_1^2 &= q_1/q_2, & \psi_1^3 &= q_1/q_3, \\ \psi_2^1 &= q_1, & \psi_2^3 &= q_3, \\ \psi_3^1 &= q_1, & \psi_3^2 &= q_2. \end{aligned} \quad (4.24)$$

These invariants have linearly independent gradients in  $\mathbb{R}^3$ .

Because the second and third eigenpair are linearly degenerate,  $\mathbf{q}_0$  and  $\mathbf{q}_1$  are connected via a single intermediate state  $\tilde{\mathbf{q}}_{1/2}$ . This intermediate state is immediately obtained from (3.8), (4.24):

$$\tilde{\mathbf{q}}_{1/2} = \begin{cases} \frac{u_1}{u_0} \mathbf{q}_0, & u_0 \neq 0, \\ \mathbf{0}, & \text{otherwise.} \end{cases} \quad (4.25)$$

#### 4.4 Two-Phase Flow

In this section we derive the intermediate states for the Euler equations for an immiscible, compressible two-phase flow. The phases are supposed to be separated by a moving interface, which is described by the time dependent set  $\mathcal{I}(t) = \{\mathbf{x} \in \mathbb{R}^3 \mid \theta(\mathbf{x}, t) = 0\}$ . Furthermore, we assume  $\theta(\mathbf{x}, t)$  to be negative in one phase and positive in the other. As a result of the immiscibility of the phases, the following kinematic condition applies:

$$\partial_t \theta + \mathbf{u} \cdot \vec{\nabla} \theta = 0, \quad (4.26)$$

where  $\mathbf{u} \in \mathbb{R}^3$  again denotes the fluid velocity. Employing the continuity equation for compressible fluids, we can restate kinematic condition (4.26) in conservation form:

$$\partial_t \rho \theta + \vec{\nabla} \cdot \rho \theta \mathbf{u} = \rho \left( \partial_t \theta + \mathbf{u} \cdot \vec{\nabla} \theta \right) + \theta \left( \partial_t \rho + \vec{\nabla} \cdot \rho \mathbf{u} \right), \quad (4.27)$$

The first term in parentheses vanishes due to (4.26), the second due to continuity. Hence,  $\rho \theta$  is a conserved quantity. Suppose that throughout the entire fluid volume the pressure is related to the density via an equation of state of the form  $p = p(\theta, \rho)$ . Then, again using  $u, v, w$  to designate the velocity components relative to a Cartesian coordinate system and ignoring spatial derivatives in  $y$  and  $z$  direction, we retrieve (2.1a), with  $\mathbf{q} = (\rho u, \rho v, \rho w, \rho \theta, \rho)^T$  and

$$\mathbf{f}(\mathbf{q}) = \left( q_1^2/q_5 + p(q_4/q_5, q_5), q_1 q_2/q_5, q_1 q_3/q_5, q_1 q_4/q_5, q_1 \right)^T. \quad (4.28)$$

Equations (2.1a), (4.28) constitute the one dimensional Euler equations for an immiscible, compressible two phase flow.

Our first objective now is to derive Riemann invariants for (2.1a), (4.28). We define  $c_1 = c_1(\theta, \rho) = \sqrt{\partial_{\theta} p}$  and  $c_2 = c_2(\theta, \rho) = \sqrt{\partial_{\rho} p}$ . Then, the Jacobian of (4.28) reads:

$$\mathbf{A}(\mathbf{q}) = \partial_{\mathbf{q}} \mathbf{f} = \begin{pmatrix} 2q_1/q_5 & 0 & 0 & c_1^2/q_5 & -q_1^2/q_5^2 - c_1^2 q_4/q_5^2 + c_2^2 \\ q_2/q_5 & q_1/q_5 & 0 & 0 & -q_2 q_1/q_5^2 \\ q_3/q_5 & 0 & q_1/q_5 & 0 & -q_3 q_1/q_5^2 \\ q_4/q_5 & 0 & 0 & q_1/q_5 & -q_4 q_1/q_5^2 \\ 1 & 0 & 0 & 0 & 0 \end{pmatrix}. \quad (4.29)$$

The eigenvalues and eigenvectors of  $\mathbf{A}(\mathbf{q})$  are

$$\lambda_1 = q_1/q_5 - c_2, \lambda_{2,3,4} = q_1/q_5, \lambda_5 = q_1/q_5 + c_2, \quad (4.30)$$

and

$$\begin{aligned} \mathbf{r}_1 &= (q_1/q_5 - c_2, q_2/q_5, q_3/q_5, q_4/q_5, 1)^T, \\ \mathbf{r}_2 &= (0, 1, 0, 0, 0)^T, \\ \mathbf{r}_3 &= (0, 0, 1, 0, 0)^T, \\ \mathbf{r}_4 &= (q_1 c_1^2, 0, 0, -c_2^2 q_5^2 + c_1^2 q_4, q_5 c_1^2)^T, \\ \mathbf{r}_5 &= (q_1/q_5 + c_2, q_2/q_5, q_3/q_5, q_4/q_5, 1)^T. \end{aligned} \quad (4.31)$$

The eigenvalue  $\lambda_k$  and the eigenvector  $\mathbf{r}_k$  are genuinely nonlinear for  $k = 1, 5$  and linearly degenerate for  $k = 2, 3, 4$ . Riemann invariants can now be obtained by solving (2.8), (4.31):

$$\begin{aligned} \psi_1^2 &= q_1/q_5 + \phi, & \psi_1^3 &= q_2/q_5, & \psi_1^4 &= q_3/q_5, & \psi_1^5 &= q_4/q_5, \\ \psi_2^1 &= q_1/q_5, & \psi_2^3 &= q_3, & \psi_2^4 &= q_4, & \psi_2^5 &= p, \\ \psi_3^1 &= q_1/q_5, & \psi_3^2 &= q_2, & \psi_3^4 &= q_4, & \psi_3^5 &= p, \\ \psi_4^1 &= q_1/q_5, & \psi_4^2 &= q_2, & \psi_4^3 &= q_3, & \psi_4^5 &= p, \\ \psi_5^1 &= q_1/q_5 - \phi, & \psi_5^2 &= q_2/q_5, & \psi_5^3 &= q_3/q_5, & \psi_5^4 &= q_4/q_5, \end{aligned} \quad (4.32a)$$

with  $p = p(\theta, \rho)$  and  $\phi = \phi(\theta, \rho)$  defined by

$$\phi(\theta, \rho) = \int_0^{\rho} \frac{c_2(\theta, \eta)}{\eta} d\eta. \quad (4.32b)$$

Observe that  $\theta$  is a  $k$ -Riemann invariant for  $k \in \{1, 5\}$ . Hence, it may be inferred that the phase transition is a contact discontinuity. Moreover, because both  $u$  and  $p$  are  $k$ -Riemann invariants for  $k \in \{2, 3, 4\}$ , the pressure and the normal velocity component are continuous across the interface.

The intermediate states can now be obtained from (3.8), (4.32). Because the linearly degenerate eigenvalue  $q_1/q_5$  has algebraic multiplicity 3, only two intermediate states have to be distinguished. Trivially,

$$\begin{pmatrix} \tilde{v}_{1/3} \\ \tilde{w}_{1/3} \\ \tilde{\theta}_{1/3} \end{pmatrix} = \begin{pmatrix} v_0 \\ w_0 \\ \theta_0 \end{pmatrix}, \quad \begin{pmatrix} \tilde{v}_{2/3} \\ \tilde{w}_{2/3} \\ \tilde{\theta}_{2/3} \end{pmatrix} = \begin{pmatrix} v_1 \\ w_1 \\ \theta_1 \end{pmatrix}, \quad (4.33)$$

and  $\tilde{u}_{1/3} = \tilde{u}_{2/3} \equiv \tilde{u}_{1/2}$ . Then,  $\tilde{\rho}_{1/3}$  and  $\tilde{\rho}_{2/3}$  are determined by

$$\begin{aligned} \phi(\theta_0, \tilde{\rho}_{1/3}) + \phi(\theta_1, \tilde{\rho}_{2/3}) &= u_0 - u_1 + \phi(\theta_0, \rho_0) + \phi(\theta_1, \rho_1), \\ p(\theta_0, \tilde{\rho}_{1/3}) &= p(\theta_1, \tilde{\rho}_{2/3}). \end{aligned} \quad (4.34)$$

We refrain from a further reduction of these expressions and suffice by stating that once the intermediate densities have been obtained,  $\tilde{u}_{1/2}$  follows by straightforward computation.

## 5. CONCLUSIONS

In spite of the absence of shock waves in most hydrodynamic applications, sufficient reason remains to employ Godunov-type schemes in this field. The shock capturing ability of these schemes renders them notably useful in the case of two-phase flow. In the present work we developed an Osher-type Riemann solver and we investigated several of its applications in the field of hydrodynamics. First, the Riemann problem was examined. Subsequently, Osher's approximate Riemann solver was discussed. It was shown that this scheme employs a rarefaction-waves-only approximate Riemann solution and that this approximation is accurate even in the presence of (weak) shocks. Then, it was demonstrated that the centered flux approximation obtained by means of Osher's scheme is not necessarily accurate and, therefore, a modified scheme was proposed. Finally, details were presented for several applications in hydrodynamics.



## Table of Contents

1	Introduction . . . . .	1
2	Riemann Problem . . . . .	2
	2.1 Preliminary Analysis . . . . .	2
	2.2 Solution . . . . .	3
3	Approximate Riemann Solution . . . . .	5
	3.1 Osher's scheme . . . . .	6
	3.2 Accuracy . . . . .	7
	3.3 Modified Scheme . . . . .	7
4	Applications in Hydrodynamics . . . . .	8
	4.1 Compressible fluid . . . . .	8
	4.2 Artificially compressible fluid . . . . .	10
	4.3 Incompressible fluid . . . . .	11
	4.4 Two-Phase Flow . . . . .	12
5	Conclusions . . . . .	14
<b>References</b>		<b>16</b>

## References

- [CHMO96] Y.C. Chang, T.Y. Hou, B. Merriman, and S. Osher. A level set formulation of Eulerian interface capturing methods for incompressible fluid flows. *J. Comput. Phys.*, 124:449–464, 1996.
- [God59] S.K. Godunov. Finite difference method for numerical computation of discontinuous solutions of the equations of fluid dynamics. *Mat. Sbornik*, 47:271–306, 1959. (In Russian).
- [KP97] F.J. Kelecy and R.H. Pletcher. The development of a free surface capturing approach for multidimensional free surface flows in closed containers. *J. Comput. Phys.*, 138:939–980, 1997.
- [Lax57] P.D. Lax. Hyperbolic systems of conservation laws II. *Comm. Pure Appl. Math.*, 10:537–566, 1957.
- [Lax73] P.D. Lax. Hyperbolic systems of conservation laws and the mathematical theory of shock waves. volume 11 of *Regional Conference Series in Applied Mathematics*. SIAM Publications, Philadelphia, 1973.
- [Lee82] B. van Leer. Flux vector splitting for the Euler equations. In E. Krause, editor, *Proc. 8th Int. Conf. on Numerical Methods in Fluid Dynamics*, volume 170 of *Lecture Notes in Physics*, pages 507–512. Springer, Berlin, 1982, Aachen 1982.
- [Lee84] B. van Leer. On the relation between the upwind-differencing schemes of Godunov, Engquist-Osher and Roe. *SIAM J. Sci. Stat. Comput.*, 5:1–20, 1984.
- [LeV90] R.J. LeVeque. *Numerical Methods for Conservation Laws*. Lectures in Mathematics. Birkhäuser, Basel, 1990.
- [MOS92] W. Mulder, S. Osher, and J.A. Sethian. Computing interface motion in compressible gas dynamics. *J. Comput. Phys.*, 100:209–228, 1992.
- [OC83] S. Osher and S. Chakravarthy. Upwind schemes and boundary conditions with applications to Euler equations in general geometries. *J. Comput. Phys.*, 50:447–481, 1983.
- [OS82] S. Osher and F. Solomon. Upwind difference schemes for hyperbolic conservation laws. *Math. Comput.*, 38:339–374, 1982.
- [Roe81] P.L. Roe. Approximate Riemann solvers, parameter vectors, and difference schemes. *J. Comput. Phys.*, 43:357–372, 1981.

- [Smo83] J. Smoller. *Shock Waves and Reaction-Diffusion Equations*. Springer Verlag, Berlin, 1983.
- [Spe87] S. Spekreijse. Multigrid solution of monotone second-order discretizations of hyperbolic conservation laws. *Math. Comput.*, 49:135–155, 1987.
- [SW81] J.L. Steger and R.F. Warming. Flux vector splitting of the inviscid gasdynamic equations with application to finite-difference methods. *J. Comput. Phys.*, 40:263–293, 1981.
- [Swe84] P.K. Sweby. High resolution schemes using flux limiters for hyperbolic conservation laws. *SIAM J. Numer. Anal.*, 21:995–1011, 1984.